

June 18, 2025

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

DOC ID# 600338

Bifenthrin GROUP 3A INSECTICIDE



BIFENTHRIN L/P Insecticide Granules

For use to control Ants (including Imported Fire Ants), Chinch Bugs, Fleas, Mole Crickets, Ticks and other insect pests on lawns in landscaped areas and perimeters around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Use areas include but are not limited to: lawns and around flower gardens, flower beds, around/along perimeters and foundations of houses and other structures, around mulch beds, along fence lines, and around decks and decking.

For outdoor use only.

Active Ingredient: Bifenthrin* 0.2%
Other Ingredient: 99.8%
Total: 100.0%
*Cis isomers 97% minimum, trans isomers 3% maximum.

FIRST AID
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® (866) 897-8050 for emergency medical treatment information.

Note to Physician - This product is a pyrethroid. If large amounts have been ingested, milk cream and other digestible fats and oils may increase absorption and should be avoided.

KEEP OUT OF REACH OF CHILDREN
CAUTION
(See other panels for additional precautionary information)

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

NET CONTENTS: 25 Lbs.

EPA Reg. No. 53883-124 (A) EPA Est. No. 5905-GA-01
(B) EPA Est. No. 88746-GA-01
First letter of lot# corresponds to EPA Est. # where product was produced



Manufactured by: Control Solutions Inc.
5903 Genoa-Red Bluff, Pasadena, TX 77507
A member of Adama Consumer and Professional Solutions
EPA 072121

WARRANTY STATEMENT
Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and that it meets the performance claims stated on the label. Control Solutions, Inc. does not warrant that the product will control all insects or mites listed on the label, or that it will control all insects or mites listed on the label under all conditions. Control Solutions, Inc. does not warrant that the product will control all insects or mites listed on the label under all conditions.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
Pesticide Disposal: Pesticide wastes are acutely hazardous.
Sprayer (all models) 2.6
Gandy Models 1.2
LESCO ROTARY 3
Scott's R-8-1 Rotary 3
Gandy Models 1.2
Equipment Disposal 1.2
EPA 800-425-5522 or www.control-solutions.com

Table with columns: Equipment, Spreader Settings, and Application Rate. Includes rows for various equipment like Gandy 5500 Series, Leeco Rotary, and Scott's R-8-1 Rotary.

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.
Do not apply directly to lawns or ornamentals. Do not apply to areas where the product will be applied to lawns or ornamentals.

Table with columns: Application Rate, Spreader Settings, and Equipment. Includes rows for various equipment like Gandy 5500 Series, Leeco Rotary, and Scott's R-8-1 Rotary.

ENVIRONMENTAL HAZARDS
This pesticide is extremely toxic to fish and aquatic life. Do not apply to water bodies, streams, rivers, or other aquatic environments.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Table with columns: Pest, Application Rate, and Active Ingredients lbs. Includes rows for Ants, Chinch Bugs, Fleas, Mole Crickets, and Ticks.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Signs of Renewed Insect Activity
If signs of renewed insect activity are observed, re-treatment may be necessary.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

LAWN & PERIMETER APPLICATIONS
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.
Do not apply directly to lawns or ornamentals. Do not apply to areas where the product will be applied to lawns or ornamentals.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.

Application Rates for Lawn and Perimeter
Under normal conditions, the treatment rates listed in the table below will control the pests listed in the table.
Apply in cool weather (65 to 80°F) in early morning or late evening hours to achieve the best results.